

ASREC Graduate Program 2018
Technical Note: The Replicator Dynamic

Jean-Paul Carvalho
Department of Economics
& Institute for Mathematical Behavioral Sciences
University of California, Irvine

22 March 2018
IRES, Chapman University

Evolutionary Game Theory

- ▶ Game theory was initially developed by mathematicians and economists.
- ▶ Evolutionary biologists adapted these techniques/concepts in developing evolutionary game theory—see for e.g. the pioneering work of British biologist John Maynard Smith. EGT was later imported back into economics.
- ▶ Owing to this intellectual history, we shall derive the **replicator dynamic** from a basic biological approach to evolution in which payoffs represent reproduction rates.
- ▶ Note that the replicator dynamic can also be generated at the macro level by imitation of more successful agents.

The Biological Approach

Ingredients:

1. Inheritance:

- ▶ Players are *programmed* with a strategy. (Players are essentially strategies.)

2. Selection:

- ▶ Strategies that do better, given what everyone else is doing, proliferate.
- ▶ In particular, payoffs are interpreted as *reproduction rates* of strategies.
- ▶ Extends Darwin's notion of survival of the fittest from an exogenous environment to an interactive setting.

3. Mutation:

- ▶ Equilibrium states can be perturbed by random shocks.
- ▶ To be *stable*, an equilibrium must be resistant to invasion by "mutant strategies".

The Strategic Context

Players

- ▶ The population consists of a continuum of agents.

Strategies

- ▶ The set of (pure) strategies is $S = \{1, \dots, n\}$, with typical members i, j and s .
- ▶ The mass of agents programmed with strategy i is m_i , where $\sum_{i=1}^n m_i = m$.
- ▶ Let $x_i = \frac{m_i}{m}$ denote the proportion of players choosing strategy $i \in S$.

The Strategic Context

Population States

- ▶ The set of population states (or strategy distributions) is $X = \{x \in \mathbb{R}_+^n : \sum_{i \in S} x_i = 1\}$.
- ▶ X is the unit simplex in \mathbb{R}^n .
- ▶ The set of vertices of X are the pure population states—those in which all agents choose the same strategy.
- ▶ These are the standard basis vectors in \mathbb{R}^n :

$$e_1 = (1, 0, 0, \dots), e_2 = (0, 1, 0, \dots), e_3 = (0, 0, 1, \dots), \dots$$

Payoffs

- ▶ A *continuous* payoff function $F : X \rightarrow \mathbb{R}^n$ assigns to each population state a vector of payoffs, consisting of a real number for each strategy.
- ▶ $F_i : X \rightarrow \mathbb{R}$ denotes the payoff function for strategy i .

Some Useful Facts

Consider the expected payoff to strategy i if i is matched with another strategy drawn uniformly at random from the population to play the following *two-player* game:

	1	2	...	n
i	$u(i, 1)$	$u(i, 2)$...	$u(i, n)$

The expected payoff to strategy i in state x is:

$$\begin{aligned}F_i(x) &= x_1 u(i, 1) + x_2 u(i, 2) \dots + x_n u(i, n) \\ &= \sum_{j=1}^n x_j u(i, j) \\ &= \sum_{j=1}^n x_j F_i(e_j).\end{aligned}$$

Some Useful Facts

The **average payoff** in the population is:

$$\begin{aligned}\bar{F}(x) &= x_1F_1(x) + x_2F_2(x) \dots + x_nF_n(x) \\ &= \sum_{i=1}^n x_iF_i(x).\end{aligned}$$

Note: this is the same as the payoff from playing the mixed strategy x against itself.

The Replicator Dynamic

- ▶ Suppose that payoffs represent *fitness* (rates of reproduction) and reproduction takes place in continuous time.
- ▶ This yields a continuous-time evolutionary dynamic called the **replicator dynamic** (Taylor and Jonker 1978).
- ▶ The replicators here are pure strategies that are copied without error from parent to child.
 - ▶ As the population state x changes, so do the payoffs and thereby the fitness of each strategy.
- ▶ The replicator dynamic is formalized as a (deterministic) system of ordinary differential equations without mutation.
- ▶ We will (at first) analyze the role of mutations indirectly by examining the local stability of various limit states of the replicator dynamic.

The Replicator Dynamic

- ▶ Let the rate of growth of strategy i be:

$$\frac{\dot{m}_i}{m_i} = [\beta - \delta + F_i(x)],$$

where β and δ are “background” birth and death rates (which are independent of payoffs).

- ▶ This is the interpretation of payoffs as fitness (reproduction rates) in biological models of evolution.

Derivation

What is the rate of growth of x_i , the *population share* of strategy i ?

By definition:

$$\begin{aligned}x_i &= \frac{m_i}{m} \\ \ln(x_i) &= \ln(m_i) - \ln(m) \\ \frac{\dot{x}_i}{x_i} &= \frac{\dot{m}_i}{m_i} - \frac{\dot{m}}{m} \\ &= [\beta - \delta + F_i(x)] - \sum_{j=1}^n x_j [\beta - \delta + F_j(x)] \\ &= F_i(x) - \bar{F}(x).\end{aligned}$$

That is, the growth rate of a strategy equals the excess of its payoff over the average payoff.

Some Properties of the Replicator Dynamic

The following results are immediate:

- ▶ Those subpopulations that are associated with better than average payoffs grow and *vice versa*.
- ▶ The subpopulations associated with pure best replies to the current population state $x \in X$ have the highest growth rate.
- ▶ $\dot{x}_i = x_i[F_i(x) - \bar{F}(x)]$, so that if $m_i = 0$ at T , then $m_i = 0$ for all $t > T$.

Example: Pure Coordination

Pure Coordination

	1	2
1	1 1	0 0
2	0 0	2 2

$$\frac{\dot{x}_1}{x_1} = (1 - x_1)(3x_1 - 2).$$

Therefore, $\frac{\dot{x}_1}{x_1} > 0$ iff $3x_1 > 2$ or $x_1 > \frac{2}{3}$.

Example: Impure Coordination

Stag Hunt

	1	2
1	2, 2	0, 0
2	0, 2	3, 3

$$\frac{\dot{x}_1}{x_1} = (1 - x_1)(3x_1 - 1).$$

Therefore, $\frac{\dot{x}_1}{x_1} > 0$ iff $3x_1 > 1$ or $x_1 > \frac{1}{3}$.

Example: Anti-Coordination

Hawk Dove

	1	2
1	-2	0
2	4	0

$$\frac{\dot{x}_1}{x_1} = (1 - x_1)(4 - 6x_1).$$

Therefore, $\frac{\dot{x}_1}{x_1} > 0$ iff $6x_1 < 4$ or $x_1 < \frac{2}{3}$.